

Study of Ethno-Botanical Flora of Dakingari, Kebbi State Nigeria, West Tropical Africa

Dharmendra Singh ¹

Abstract

The present research is an attempt to study the ethno-botanical flora of Dakingari, Kebbi State, Nigeria. The study was conducted from July 2014 to June 2015 villages of Dakingari. The used of questionnaire method was used for obtaining information of the indigenous knowledge of plants that are used traditionally. The results were spreads in about 31 families, 45 genera and 53 species of angiospermic families. They are Amaranthaceae 2 species, Anacardiaceae 3species, Annonaceae 1 species, Asclepiadaceae 2 species, Asteraceae (Compositae) 2 species, Balanitaceae 1 species, Bombacaceae 1 species, Caesalpinaceae 7 species, Caricaceae 1 species, Capparaceae 1 species, Combretaceae 4 species, Convolvulaceae 1 species, Cucurbitaceae 1 species, Ebenaceae 1 species, Euphorbiaceae 4 species, Lilliacae 1 species, Malvaceae 2 species, Meliaceae 1 species, Mimosaceae 2 species, Musaceae 1 species, Moringaceae 1 species, Mrytaceae 2 species, Nymphaeaceae 1 species, Olacaceae 1 species, Pediliaceae 2 species, Poaceae 2 species, Rubiaceae 2 species, Sapindaceae 1 species, Sapotaceae 1 species, Sterculiaceae 1 species and Verbanaceae 1 species.

Keywords: *Ethno-botany, Dakingari, Flora, Kebbi State.*

¹ Department of Biological Sciences Kebbi State University of Science and Technology Aliero, Kebbi State PMB-1144, Nigeria (singhdharmendra12@yahoo.com)

Article history: Received February 10, 2016; Revised March 14, 2016; Approved April 5, 2016

Bi-Monthly Double-Blind Peer Reviewed Refereed Open Access International E-Journal - Included in the International Serial Directories Indexed & Listed at: Thomson Reuters Researcher ID: C-6767-2016, Google Scholar as well as in Directory of Abstract Indexing for Journals, U.S.A.

Scopus ID: 9F46766090560E8E; Impact Factor Evaluation [SJIF 2015 = 3.605]
International Research Journal of Engineering, IT and Scientific Research

I. Introduction

Though the term “ethno botany; was not coined until 1895 by the US botanist John William harsh berger, the history of the field begins long before that Pythagoreans, which originated in 500 BC included refusal to the beans, perhaps because of the human relationship of beans through matter (Baruah and Kalita,2007). Theophrastus the father of botany wrote of plant and people usage of them in his works. In A.D. 77, the Greek surgeon Pedanius Dioscorides published De Material Medical, which was a catalog of about plant in the Mediterranean). It also included information on how the plants were used, especially for medicinal purpose. This illustrated herbal publication contained information on how and when each plant was gathered, whether or not it was poisonous, its actual use, and whether or not it was edible (it even provided recipes). Dioscorides stressed the economic potential of plants. (Sarmah, 2000).

The uses of natural herbal drugs whether traditional or modern, have originated directly or indirectly from folklore, and rituals known as ethno medico-botany. The use of herbal medicine reflects the long history of human interaction with the environment (Sarmah, 2010).

During the mediaeval period ethno botanical studies were commonly found connected with monasticism. Notable at this time was Hildegard von Bingen. However most ethno botanical knowledge was kept in garden such as physic garden attached to hospital and religious building. It was thought of in practical, use term for culinary and medical purpose and the ethnographic element was not studied as modern anthropologist might approach ethno botany today.

Beginning of the 20thcentaury, the field of ethno botany experienced a shift from the raw compilation of data to a greater methodological and conceptual reorientation. This is also the beginning of academic ethno-botany. The so-called father of this discipline is Richard Evans Schulte’s even though he did not actually coin the term Ethno-botany’. Today the field of ethno botany requires a variety of skills; botanical training for the identification and preservation of plant specimen; anthropological training to understand the cultural concept around the perception of plants; linguistic training, at least enough to transcribe local terms and understand native morphology, syntax, and semantics.

The publication of 'Flora of West Africa' by (Hutchinson and Dalziel 1963) was followed by two national floras by (Stanfield, 1970) for Grasses of Nigeria (Lowe and Stanfield 1974) for Flora of Nigeria (Sedges). Later on some aspect of studies in floridiversity of Kebbi state, Nigeria by (Singh *et al.*, 2010), towards making a flora of Kebbi state, Nigeria by (Singh and Misra 2013 a ,b). Morpho-taxonomic and Ethno – medicinal flora of Kebbi state, Nigeria by (Singh and Abubakar 2011a,b,c, Singh 2015).

II. Material and Methods

2.1 Study Area

The study area Dakingari is located at approximately at latitudes $11^{\circ} 38' 51'$ North and $4^{\circ} 3' 51'$ East It is bounded the North by Bunza local government area, East by Kalgo local government area and Maiyama local government area, South east by Bagudo local government area, West by Dandi local government area. It is the fourth largest local government located in the central part of Kebbi state..

2.2 Climatic Condition

The climate conditions of Dakingari enjoy tropical type of climatic condition generally characterized by two different seasons that is wet and dry season. The rainfall began by the month of April with the heaviest rainfall in the month of July-to August with annual rainfalls between 500 mm to 1300 mm. The dry season is characterized with few or absence of vegetation cover and non- incidence of rainfall. It is a period of low agricultural activities mainly restricted to Fadama. This season is also made up of two period which are cold harmattan, recorded in the month of November to January and hot harmattan recorded between the month of February.



Fig. Map of Dakingari area

2.3 Plants collection

The study was conducted from July 2014 to June 2015 villages of Dakingari. The used of questionnaire method was used for obtaining information of the indigenous knowledge of plants that are used traditionally. The plants specimens collected from different localities and habitats in every stage of their growth and reproduction. A complete specimen possesses all part of plants including root system. The plants specimen should be usually in flowering stage. The specimen collected most be tagged immediately and record about localities most also be noted (Jain and Rao 1978, Okoli 2002, Tucker and Calabrese 2005).

2.4 Pressing and drying of plants

The plants pressed in between the sheet of blotting papers or old newspapers. One plant be arranged on one sheet in a manner that there should be no overlapping of plants

(Jain and Rao 1978). The blotting papers with plant specimen were placed in field press for about 24 to 48 hours. The press is then opened blotting should be change and rearrange the plants properly. After it press should be again closed and again after 2 or 3 days change the blotting and dry plants in sunlight or artificial heat (Jain and Rao 1978).

2.5 Mounting and labeling of the specimens

After drying, the specimen mounted for permanent record on sheet called mounting papers on herbarium sheets. The specimen are mounted to the sheet with the help of fevicol or quick drying liquid paste after mounting the specimens on the herbarium sheet labeled. A label is pasted or printed on lower right-hand corner, compressed in plant press to prevent them from drying and this press herbarium were house in the herbarium of Department of Biological Sciences, Kebbi State University of Science and Technology Aliero, Nigeria for record and references

III. Justification of Research

Ethno-botany is the study of how people of a particular culture and region make use of indigenous plants. In the study, ethno-botanists gather data from the people with the view of understanding the ways by which they use plants for medicine, food, construction materials and as tools. Before the advent of western medicine, indigenous people worldwide have been taking care of their health needs using plants. The African continent has a long history of the use of plants, in some African countries up to 80% of the rural population rely on medicinal plants as a source of remedies.

IV. Enumeration of Taxa

The present investigation enumerates the taxa during the study of ethno-botanical flora of Dakingari, Kebbi State, Nigeria. The study was conducted from July 2014 to June 2015 villages of Dakingari. After a thorough study on 250 field numbers, the identified taxa have been found to be spread about 53 species and 45 genera of 31 families, these have been arranged in an alphabetical order, and the same pattern has been followed in arranging lower order taxa under each family. Each taxa have been discussed their ethno-botanical uses.

1) Amaranthaceae

(1) Botanical name: *Amaranthus viridis* L.

Local name: Alayyahu

Ethno-botanical uses: The leaves are edible and also squeezed to extract its water content that are used in curing eye disease by dropping few drops water twice a day.

Dakingari, D.Singh: 36.

(2) Botanical name: *Amaranthus spine* L.

Local name: Rukkubu

Ethno-medicinal uses: The plant is edible the leave and tender stems are eating, cooking.

Dakingari, D.Singh: 28.

2) Anacardiaceae

(3) Botanical name: *Anacardium occidentale* L.

Local name: Kashu

Ethno-botanical uses: The fruit is edible and chewed to treat vomiting, nausea and relief for liver diseases. Bark is dried and grind to powder, add to pap and drunk for cough, jaundice and dysentery. Leaves are boiled and take for measles.

Dakingari, D.Singh: 39.

(4) Botanical name: *Mangifera indica* L.

Local name: Mangwaro

Ethno-botanical uses: Fruit is edible. Leaves are prepared through decoction for remedy of diarrhea, fever, chest pain, and concoction of mango leaf with other plant such as guava, orange, is taken after child birth for body strength.

Dakingari, D.Singh: 01.

(5) Botanical name: *Sclerocarya birrea* (A. rich) hochst.

Local name: Lodaloda

Ethno-botanical uses: Ripe fruit are edible, leaves is used for treat sore throat. Bark combined with that of *Parkia biglobosa* for the treatment of syphilis and measles.

Runtowa, D.Singh: 53.

3) Annonaceae

(6) Botanical name: *Annona senegalensis* L.

Local name: Gwanda Daji

Ethno-botanical uses: The ripe fruit is edible and the whole part of the plant is use as pain relieve and also to treat gastro-Intestinal disorder.

Uroba, D.Singh: 31

4) **Asclepiadaceae**

(7) Botanical name: *Calotropis procera* (L) R.Br.

Local name: Tunfafiya

Ethno-botanical uses: The plant is medicinal the bark is grinded in to powder and use in curing some common diseases such as inflammatory disease, fever, rheumatism, tuberculosis, the latex contents also rub in the body for skin diseases such as eczema.

Dakingari, D. Singh: 24

(8) Botanical name: *Sarcostema viminale* Smith.

Local name: Bakinmutum

Ethno-botanical uses: The root are boiled and drunk for the treatment of dysentery, stomach pain.

Uroba, D.Singh: 27

5) **Asteraceae**

(9) Botanical name: *Vernonia amygdalina* L.

Local name: Shuwaka

Ethno-botanical uses: The leaves are used to prepared soup. It also impounded squeezed add with milk and drunk for heart disease and increase milk production in women.

Dakingari and Lambu, D.Singh: 10

6) **Balanitaceae**

(10) Botanical name: *Balanite aegyptica* Delite.

Local name: Adua

Ethno-botanical uses: Leaves are dried and grind to powder, add to pap or honey to treat stomach-ache, yellow fever. Bark also soaked in water to treat dysentery.

Gandu, D.Singh: 5

7) **Bombacaceae**

(11) Botanical name: *Adansonia digitata* L.

Local name: Kuka

Ethno-botanical use: Dried leaves are grind in to powder and prepare a soup, fruit are edible bark are boiled and taken as a treatment of malaria, asthma, diarrhea, bladder diseases at least once a day not too much.

Dakingari, D.Singh: 48.

8) **Caesalpiniaceae**

(12) Botanical name: *Cassia mimosoides* L.

Local name: Baguruwar kasa

Ethno-botanical use: Its use to dysentery, headache, fever through squash method, when is grinded in to powder it treat rashes after child birth.

Dakingari, D.Singh: 35

(13) Botanical name: *Cassia tora* L.

Local name: Tafasa

Ethno-botanical uses: The cooked leaves are edible. It is also very useful in treating skin disease like ringworm, itching or body scratch, fresh leave is use externally to treat eczema and dermatomycosis, through squash method. Fruits are decocted and drink for treating fever.

Dakingari, D. Singh: 37

(14) Botanical name: *Temarindus indica* L.

Local name: Tsamiya

Ethno-botanical uses: The ripe fruit is said to treat fever, intestinal discomfort ailment in case of poisoning, infection swelling .The fruit also reduces blood sugar.

Uroba: D.Singh 21

(15) Botanical name: *Cassia kotschyana* Oliv.

Local name: Malga

Ethno-botanical uses: Root, with combination of *Guiera senegalensis* are boiled and used for steam-bath for the treatment of general body pain.

Dakingari guba, D.Singh: 40

(16) Botanical name: *Piliostigma reticulate* Sch.

Local name: Kalgo

Ethno-botanical uses: Root with combination of *Guiera senegalensis* and *Waltheria indicca* and marked, is used to treat serious diarrhea both adult and children.

Dakingari and Road site, D.Singh: 44

(17) Botanical name: *Cassia occidentalis* L.

Local name: Sanga Sanga

Ethno-botanical uses: Decoction of fresh leaf with combination of other plant such as orange, dogonyaro. Is said to treat malaria fever, typhoid fever, Rheumatism and rip pain both adult and child.

Dakingari, D.Singh: 14

(18) Botanical name: *Senna alata* L. Roxb.

Local name: Filaskon bature

Ethno-botanical uses: Leaves are boiled and drunk is said to treat malaria fever, typhoid fever and dysentery.

Dakingari, D. Singh: 17

9) Caricaceae

(19) Botanical name: *Carica papaya* L.

Local name: Gwanda

Ethno-botanical uses: The fruit is edible, leaves, seeds, are used to treat gonorrhea, Syphilis, Amoebic dysentery, through decoction method the fruit are also use in treating ringworm, malaria through grinding in to powder the fruit is also edible.

Dakingari, D.Singh; 45

10) Capparaceae

(20) Botanical name: *Cleome viscosa* L.

Local name: 'Yar Unguwa

Ethno-botanical uses: The plant is edible fresh is squeezed and place in to anus for treatment of dysentery.

Dakingari, D.Singh : 32

11) Combretaceae

(21) Botanical name: *Guiera senegalensis* (L) Briq

Local name: Sabara

Ethno-botanical uses: The plant is medicinal and is used in treatment of allergic reaction such as thrush; fresh leave is grinded into powder to rub all over the body. It is also used by woman after given birth promote their body' strength through taking bath

Wasadabiri, D.Singh: 3

(22) Botanical name: *Terminalia avicennioide* Gull.et Perrott.

Local name: Baushe

Ethno-botanical use: Bark is boiled and taken orally to treat cough, syphilis, rheumatism and joint pain.

Polytechnic campus, D.Singh: 22

(23) Botanical name: *Combretum fragans* F. Hoffm.

Local name: Jar tarauniya

Ethno-botanical uses: The leaves boiled extensively and drunk to cease blood bleeding, for syphilis, body pain and itchiness. Leaves and root for asthma, chest pain.

Uroba, D.Singh: 26.

(24) Botanical name: *Combretum lecardii* Engl and Diels.

Local name: Farartarauniya

Ethno-botanical uses: The leaves boiled extensively and drunk to stop bleeding, and also use treat syphilis, body pain and itchiness. Leaves and root for asthma, chest pain and constipation.

Uroba, D.Singh: 25

12) Convolvulaceae

(25) Botanical name: *Ipomoea repens* L.

Local name: Dumal Kada

Ethno-botanical use: The whole plant both root, stem, flower are boiled then a patient of dysentery enter the water and sit-down at least 30m. it is also used for treatment of syphilis.

Dakingari and Runtowa, D.Singh:42

13) Cucurbitaceae

(26) Botanical name: *Citrillus vulgaris* Thumb.

Local name: Kan kana

Ethno-botanical use: The fruit are edible. The rind of fruit is used to treat ulcer with combination of milk it also said to treat diabetic, hypotension.

Dakingari, D.Singh: 33

14) Ebenaceae

(27) Botanical name: *Diospyros piliformis* Hochst. ex. A.DC.

Local name: Kanya/Kaiwa

Ethno-botanical uses: The fruits are edible. The fresh fruits are grinded and filtered and a cup of the filtered extract is taken twice a day for remedy of fever, diarrhea.

Dakingari, D.Singh: 13

15) Euphorbiaceae

(28) Botanical name: *Euphorbia hirta* L.

Local name: Nonon Kurchiya

Ethno-botanical uses: The fresh leave, stem and seed are chewing in case of Asthma, bronchitis, worm infection (female) and dysentery. It is taken frequently.

Dakingari, D.Singh: 16

(29) Botanical name: *Flueggea virosa* Roxb.

Local name: Tsa

Ethno-botanical uses: The whole parts of the plants, fruit are boiled and water is added to pap and drunk for diarrhea. Bark is chewed for stomach-ache.

Dakingari and Uroba, D.Singh: 20

(30) Botanical name: *Phyllanthus pentendrous* Sch and thon.

Local name: Geron tsuntsaye

Ethno-botanical uses: The whole plant are chewed for throat sore and ear problem and also used for the stomach-ache.

Dakingari, D.Singh: 51

(31) Botanical name: *Ricinus communis* L.

Local name: Dankwasaro

Ethno-botanical uses: Fruits are pounded dissolved in water sieved and drunk to prevent being pregnancy from women.

Tungar lanta, D.Singh:52

16) Liliaceae

(32) Botanical name: *Alluum cepa* L.

Local name: Albasa

Ethno-botanical uses: The bulb, leave extract are used to treat inflammation, dental problem, throat infection by squeezing and rubbing on the affected area and the bulb is edible, and used for food.

Lambu, D.Singh: 49

17) Malvaceae

(33) Botanical name: *Hibiscus sabdariffa* L.

Local name: Zoborodosobo

Ethno-botanical uses: The back of the fruit is use as Roselle tea and is said to treat high blood pressure, cold, tooth ache, urinary tract infection, conjunctivitis it also increase the volume of blood and it edible cooking soup.

Dakingari, D.Singh 18

(34) Botanical name: *Abelmoschus esculentu* L.

Local name: Kubewa

Ethno-botanical uses: The fruit is edible cooking making soup and it increase blood that is blood builder.

Dankingari, D.Singh: 47

18) Meliaceae

(35) Botanical name: *Azadirachta indica* A. Juss.

Local name: Dogonyaro

Ethno-botanical use: The fresh leave is highly ingredient with antiseptic anti-inflammatory gingivitis i.e gum infection malaria, typhoid fever through squash method it is taken very small not much because it is has bitter taste.

Dakingari, D.Singh:43

19) Mimosaceae

(36) Botanical name: *Acacia nilotica* L willd exde.

Local name: Bagaruwa

Ethno-botanical uses: The fresh leave is squeeze and taken as a remedy of stomach pain, ripe dried fruit is grinded in to powder with combination of milk are taken orally for the treatment of ulcer, at least twice a day not much.

Dakingari, D.Singh: 7

(37) Botanical name: *Acacia erythrocalyx* (L) Willd.

Local name: Gumbi

Ethno-botanical uses: The roosts are grinded to powder for wounds. Bark is chewed for tooth decay. Leaves are boiled for headache and cough.

Dakingari, D.Singh: 12

20) Musaceae

(38) Botanical name: *Musa sapientum* L.

Local name: Ayaba

Ethno-botanical uses: The ripe fruit is edible as plantain. The leaves are boiled with combination of plant such as *Cassia occidentals* and *Psidium guajaba* are taken as remedy of malaria, typhoid fever, diarrhea and mental disorder.

Dakingari and Lambu, D.Singh: 46

21) Moringaceae

(39) Botanical name: *Moringa oleifera* L.

Local name: Zogale

Ethno-botanical uses: The fresh plants are boiled and taken the water for the treatment of paralysis, eye diseases, asthma, the leave are cooked and edible the water is also taken most especially by diabetic patient.

Dakingari and Lambu, D.Singh: 2

22) Myrtaceae

(40) Botanical name: *Psidium guayava* L.

Local name: Gwaiba

Ethno-botanical uses: The ripe fruit is edible. Leaves are boiled and drunk to treat malaria, typhoid fever, with combination of *Cassia occidentalis*, orange, mango. Steam-bath also performed morning and evening twice a day and regular taken, drinking of water.

Dakingari and Lambu, D.Singh: 9

(41) Botanical name: *Eucalyptus camaldulensis* Schlectht.

Local name: Bishyar turare

Ethno-botanical uses: The fresh leaves are combined with mango, guava, orange and boiled for steam-bath in treatment of typhoid, malaria, dysentery, cold, and cough served as body pain relievers.

Dakingari, Barbarejo and Road site, D.Singh: 15

23) Nymphaeaceae

(42) Botanical name: *Nymphaea lotus* Willd.

Local name: Bado

Ethno-botanical uses: Ripe fruits are edible. Root are also boiled and drunk as bladder problem when a person is not passing urine it will help to ease urination.

Tungar lanta and River, D.Singh: 04

24) Olacaceae

(43) Botanical name: *Xinema america* L.

Local name: Tsadan daji

Ethno-botanical uses: Ripe fruits are edible. Leaves are boiled with combinations of *Boswellia dalzielii* (hano), and *Pseudocedrela kotschy* (tuna) added potassium and drunk for inflammatory diseases, dysentery.

Uroba, D.Singh: 41

25) Pedaliaceae

(44) Botanical name: *Ceratotheca sesamoides* Endl.

Local name: Yodokarkashi

Ethno-botanical uses: The fresh leaves are squash and rub on the skin as a treatment of skin infection, leaves are used by women to treat lice problems through washing their hair with the impounded fresh leaves.

Dakingari and Gandu, D.Singh: 19.

(45) Botanical name: *Sesamum indicum* L.

Local name: Ridi

Ethno-botanical uses: The fruit is edible and used to produce oil. Leaves are squashed and rubbed on the skin as a treatment of skin infection.

Dakingari and Gandu, D.Singh: 50

26) Poaceae

(46) Botanical name: *Triticum aestivum* Lam.

Local name: Alkama

Ethno-botanical uses: It is edible and used to make food. Pap added with honeybee is said to maintain the quality of breast in women. It is also served as food of diabetic patients.

Dakingari and Lambu, D.Singh: 34.

(47) Botanical name: *Eragrostis tremala* Hotch Ex.

Local name: Burburwa

Ethno-botanical uses: The root is chewed for treatment of throat pain

Dakingari and Gandu, D.Singh: 29

27) Rubiaceae

(48) Botanical name: *Sarcocephalus latifolius* Smith.

Local name: Tafashiya/tuwonbiri

Ethno-botanical uses: The ripe fruit is edible. Leaves are boiled and drunk for the treatment of inflammation, dysentery. Leaves powdered are also added to pap for the treatment of many diseases.

Dakingari and Uroba, D.Singh: 30

(49) Botanical name: *Mitracapus hirtus* L. DC.

Local name: Harwatsi

Ethno-botanical use: The fresh plants are used in treatment of eczema through which plant is squeezed and rubbed in the affected area.

Dakingari and Gandu, D.Singh: 6

28) Sapindaceae

(50) Botanical name: *Paullinia pinnata* L.

Local name: Farce biyar

Ethno-botanical uses: Whole plants are boiled and drunk for the fasts walking and aid teeth germination in infant children.

Dakingari and Uroba, D.Singh: 38

29) Sapotaceae

(51) Botanical name: *Vitellaria paradoxa* C.F.Gaertn.

Local name: Kade

Ethno-botanical uses: The fruit are edible and uses to produces oil (man kade) fresh leaves are boiled and taken as treatment of stomach ache, rheumatic ache, diarrhea, nausea, fruit ointment are used on skin infection, and apply on head of hair growth.

Dakingari and Uroba, D.Singh: 23

30) Sterculaceae

(52) Botanical name: *Waltheria indica* L.

Local name: Damaigi

Ethno-botanical uses: The leaves are boiled and taken as a treatment of abdominal pain, diarrhea most especially for children.

Dakingari, D.Singh: 11

31) Verbanaceae

(53) Botanical name: *Vitex doniana* Shumach and Thorn.

Local name: Dunya

Ethno-botanical uses: The fruit is edible, root are boiled and take as remedy of cholera, anemia, leprosy, mouth infection.

Dakingari, D.Singh: 8

V. Result and Discussion

In this study, fifty (53) indigenous species of plants belonging to (31 families) were documented as being useful for human life. As trees have the highest number of (23 species) compared to both shrubs (12 species) and herbs (18 species) respectively. It was notice that, most of the plants have multipurpose medicinal uses and parts of each plant species being used. While some plants are edible, others are of economic importance. The plants are prepared in various ways for the medicinal purposes such as concoction,

decoction, grinding, impounding etc. of leave, stem, flower fruit, bark, and seeds. These various formulations are either, drunk sniff, smoke, bath, steaming, enema, rubbed etc. on the body for diseases management. Other plants are used as sources of food. Varieties of plants are prepared commercially as source of income in various communities. *Sesamum indica* and *Vitellallria paradoxa* plant seeds are used as sources of oil.

VI. Conclusion

From this research work we conclude that all human being and animals almost depend on plant directly or indirectly on plant as a source of energy, food, shelter, economic, medicine, fuel, fire wood and as raw materials for industries purpose. Almost all the synthetic drugs and traditional drugs are source from plants. The ethno-botany is of great importance as it provides easier and cheaper way for the management of malnourishment ailment within the society.

VII. Acknowledgement

Author is grateful to the Tertiary Education Trust Fund Abuja, Nigeria for financial support and also gratefully acknowledge to the working facilities provided by the Department of Biological Sciences, Kebbi State University of Science and Technology Aliero, Kebbi State, Nigeria

References

- [1] Baruah S. and Kalita C.(2007). Forest and Savannah 4th edition Ibadan and London. Pp. 51-55.
- [2] Hutchinson, J and Dalziel, J.M (1963). *Flora of West Tropical Africa* Vol, 1, 2, and 3. Revised by R.J.W. Crown Agent for overseas Government, U.K.
- [3] Jain, S.K and Rao, R.R (1978). *Hand book Of field and Herbarium methods*. Today and Tomorrow printers and publishers. New Delhi. Indian. Pp. 33-58.
- [4] Lowe, J. and Stanfield, D.P (1974). *Flora of Nigeria (Sedges)*. University of Ibadan press, Ibadan Nigeria.
- [5] Okoli, B.E. (2002). *Field and Herbarium Techniques*. Mbeyi Publishers Port Harcourt, Nigeria.Pp.15-20.
- [6] Sarmah C.O.(2000). A global assessment of enderonism and species richness across Island and Mainland regions. P.N.A.(106). Pp. 9322-9327.
- [7] Sarmah C.O.(2010). Research and collection of herbarium Department of Botany.National herbarium Pp. 119-124.

- [8] Singh, D. Misra, B. K. & Abubakar, A. (2010). Some aspects of the studies in floriversity of Kebbi State, Nigeria, West Tropical Africa-I. *Biological and Environmental Sciences Journal for the Tropics* 10(4) 212-218 Nigeria.
- [9] Singh, D. & Abubakar, A. (2011 a). Morpho-Taxonomic and Ethno-Medicinal Aspects of Flora in Kebbi State, North-Western Nigeria. *Biological and Environmental Sciences Journal or the Tropics* 8(2) 198-205 Nigeria.
- [10] Singh, D. & Abubakar, A. (2011 b). Study of Weed Flora of Kebbi State, Nigeria, West Tropical Africa. *Journal of Biological Sciences and Bioconservation* Vol. 2 89-96.
- [11] Singh, D. & Abubakar, A. (2011 c). Morpho-Taxonomic and Ecological study of *Aloe vera* Tourn. EX L. species found in Kebbi State, Nigeria, West Tropical Africa. *Journal of Phytological Research* 24 (2) 237-239.
- [12] Singh, D & Misra, B.K. (2013 a). Towards Making a Flora of Kebbi State, Nigeria parts: 2, West Africa. *Biological and Environmental Sciences Journal or the Tropics* Vol. 10 (4) 184-188.
- [13] Singh, D. & Abubakar, M. (2013 b). Study of the Ethno-medicinal Flora in the Aliero Local Government Area, Kebbi State, Nigeria. *National Journal of Life Sciences* 10 (2) 251-256
- [14] Stanfield, D.P (1970). *Flora of Nigeria, Grasses*. University of Ibadan press, Ibadan, Nigeria.
- [15] Singh, D.(2015). Floristic Diversity of Kebbi state, Nigeria, West Tropical Africa, Lap Lambert Academic Publishing Germany.Pp.145.
- [16] Tucker,O and Calabrese, L (2005).*The use and methods of making Herbarium /Plant specimens*, The Herb Society of America 9019 Kirtland Road, Kirtland, OH 44094. Pp. 1-20.